

A serological comparison of the populations of the Levantine Viper, *Macrovipera lebetina* (Linnaeus, 1758) in Cyprus and Southern Turkey

Bayram GÖÇMEN¹, Hüseyin ARIKAN¹,
Kerim ÇİÇEK^{1,*} and Mehmet Zülfü YILDIZ^{1,2}

1. Ege University, Faculty of Science, Department of Biology, Zoology Section, Tr 35100 Bornova, Izmir-Turkey

2. Harran University, Faculty of Art-Science, Department of Biology, Zoology Section,
Osmanbey Campus, Sanliurfa-Turkey

*Corresponding author: Kerim ÇİÇEK, E-mail: kerim.cicek@ege.edu.tr,
Tel: 0 (232) 388 40 00/1790, Fax: 0 (232) 388 10 36

Abstract: The present study compares the blood-serum proteins of populations of the Levantine viper, *Macrovipera lebetina* L. from Cyprus and southern Anatolia by polyacrylamide gel electrophoresis and densitometry analysis methods. There were discernible differences found between the two populations for the blood serum protein features compared. By considering these differences, it is concluded that the southern Anatolian population of *Macrovipera lebetina* should not be considered as the nominate subspecies *M. l. lebetina*, which lives in Cyprus.

Key Words: *Macrovipera lebetina lebetina*, *M. l. obtusa*, blood-serum proteins, Cyprus, Turkey.

Introduction

The Levantine viper, *Macrovipera lebetina*, has a rather extensive geographical range in central Asia, the Middle East and northern Africa (Algeria & Tunisia) (Nilson & Andren 1988, Nilson et al. 1988, Böhme & Wiedl 1994, Göçmen et al. 1996, Schleich et al. 1996, David et al. 1999, Atatür & Göçmen 2001).

There has been considerable confusion and disagreement among authors with regard to the taxonomic status of *M. lebetina*. Asiatic and Russian *lebetina* were attributed to the subspecies *obtusa* for a long time by

several authors (Chernov 1944, 1959, Terentjev & Chernov 1949, Bannikov et al. 1977, Başoğlu & Baran 1980, Joger 1984, Nilson & Andrén 1988, Chickin & Szczerbak 1992, Leviton et al. 1992, Baran & Atatür 1998, Tok et al. 2002), but there was no general consensus about its definition. The populations from the easternmost parts of the range of *M. lebetina*, previously regarded as belonging to the subspecies *turanica* (Chernov in Terentjev & Chernov, 1949), were described as *chernovi* by Chikin & Szczerbak (1992). Different authors argued over the validity of the other subspecies *turanica*, *euphratica*, *peilei*

and *chernovi* from Asia, and, for example, Joger (1984) considered only two valid subspecies: *lebetina* of Cyprus and *obtusa* of Asia (except Cyprus).

Although Anatolian *Macrovipera lebetina* were attributed to the subspecies *obtusa* for a long time (Baran 1976, Başoğlu & Baran 1980, Baran & Atatür 1998, Budak & Göçmen 2005), some authors (Billing & Schaetti 1984, Broodmann 1987, Ossenegg 1989, Mulder 1995) stated that the *M. lebetina* specimens from the southern parts of Anatolia closely resemble those of Cyprus (*M. l. lebetina*). Billing & Schätti (1984), Broodmann (1987) and Mulder (1995) stated that the specimens from the southern parts of Anatolia closely resemble those of Cyprus (*M. l. lebetina*) regarding head shape, color pattern, and number of ventrals. However, Arıkan *et al.* (2005) and Göçmen *et al.* (2006) comparatively examined populations from both localities regarding morphology, hemipenes and electrophoresis pattern of venom proteins. They found that significant differences between the two populations for some aspects of general morphology, hemipenial morphology and venom proteins. Therefore, they concluded that these populations are taxonomically distinct.

The present study aims to further examine the southern Turkey population from a serological point of view in order to clarify whether it should be considered as the nominate subspecies *M. l. lebetina* which lives in Cyprus.

Materials and Methods

For electrophoresis, we used a total of 8 (4 ♂♂ and 4 ♀♀) adult *Macrovipera lebetina* specimens collected in northern Cyprus (2 ♂♂ and 2 ♀♀ from Kyrenia, Nicosia and Karpas Districts) and southern Turkey (2 ♂♂ and 2 ♀♀ from Kilis, Şanlıurfa and Diyarbakır Provinces). Blood samples were obtained from the postorbital sinuses via heparinized haematocrit tubes (MacLean *et al.* 1973). For electrophoretic study, 5 µl of blood-serum proteins were separated in polyacrylamide gel disk electrophoresis using a Canalco Model 1200 disk electrophoresis apparatus. Detailed procedure was explained by Arıkan *et al.* (1998, 2005). Qualitative evaluations of the gels were done directly from the electropherograms and the densitometric curves of the separations were created by means of a Gelman ACD-15 Model 39430 densitometer scanning at 500 nm.

Results

All specimens examined were sexually mature and no obvious difference was recorded in serum protein pherograms (in densitometric curves) between the sexes in either Cypriot or southern Turkish populations. Consequently, the sexes of each population were pooled for further evaluation. The protein distribution pattern of southern Turkey and Cypriot specimens representing of the *M. lebetina* populations studied is given in Figure 1 and 2, along with their densitometric tracing curves.

In qualitative comparisons between the specimens from northern Cyprus and southern Turkey, the electropherograms clearly differed in glo-

bulin zones. In Cypriot samples (Fig. 1), the number of discernible globulin fractions or fraction groups (zones) was 11, while in samples from southern Turkey they number 10 (Fig.2). Besides, the total fraction number of the blood-serum proteins of the Cypriot *M. lebetina* specimens

comprises 12 fractions or fraction groups but 11 in the southern Turkey specimens when we include a common albumin fraction or fraction group. Additional quantitative differences (in the values of optical densities) were seen in each globulin fraction or fraction groups (fig. 1-2).




Figure no.1 Representative electropherogram (gel photograph) showing the electrophoretic separation of the blood-serum proteins of the Cypriot *Macrovipera lebetina*, together with its densitometric tracing curve. O.D.: Optical density, S: Start (junction between the spacer and separation gels).

Discussion

Many researchers have underlined the taxonomic importance of the density, speed and fraction numbers obtained from electrophoretic separation

of blood-serum proteins of amphibians and reptiles (Dessauer & Fox 1956, Chen 1967, Ferguson 1980, Arıkan et al. 1998). The qualitative differences of fractions could be caused by genetic variations, whilst

quantitative differences could reflect age, gender, environmental and physiological factors (Ferguson 1980); therefore, qualitative differences are important for taxonomic studies. In this study, the obtained data from the

electrophoresis of blood serum proteins indicate that *M. lebetina* populations from southern Anatolia and Cyprus show significant differences qualitatively.




Figure no.2 Representative electropherogram (gel photograph) showing the electrophoretic separation of the blood-serum proteins of the southern Turkish *Macrovipera lebetina*, together with its densitometric tracing curve (For further explanations, see legend to Fig. 1).

Arikan et al. (2005) and Göçmen et al. (2006) have compared the general morphology, hemipenial morphology and venom protein pattern of Cypriot and southern Turkey population and they concluded that the southern

Turkey population should not be included *M. lebetina lebetina*. In this study, we also conclude that these populations are taxonomically distinct in the viewpoint of blood serum proteins. Consequently, the popu-

lation from southern Anatolia should not be allocated to the nominate subspecies *M. l. lebetina*, which lives in Cyprus, as previously suggested by Billing & Schätti (1984), Broodmann (1987), Ossenegg (1989) and Mulder (1995).

In the relevant literature, the population of *M. lebetina* from NE Anatolia has long been referred to as *M. l. obtusa* (Mertens 1952, Eiselt & Baran 1970, Baran 1976, Nilson et al. 1988, Nilson & Andrén 1988, Joger et al. 1997, Atatür & Göçmen 2001, Tok et al. 2002). As suggested by Billing and Schätti (1984), Broodmann (1987), Ossenegg (1989) and Mulder (1995), the population existing in southern Anatolia is distinctly different at the subspecific level from the population in NE Anatolia, which should be resurrected as *M. l. euphratica* Martin, 1838 from the synonym list of the species, since its type locality was given as "shores of Euphrates".

In summary, our findings on blood serum proteins indicate that the southern Turkey population might be quite distinct from the Cypriot population at a subspecific level; and while we were unable to compare with the specimens from northeastern Turkey populations, according to the relevant literature, it is possible to suggest that they also belong to distinctly different subspecies. We thus accept, provisionally, considering the limited number of our specimens for comparison, the subspecific status of the southern Anatolian population as *Macrovipera lebetina* ssp.

References

- Arıkan, H., Atatür, M.K., Mermer, A. (1998): An investigation on the blood serum proteins of *Chalcides ocellatus* (Sauria: Scincidae) populations from Southern Anatolia. *Turkish Journal of Zoology* 22: 175-177.
- Arıkan, H., Göçmen, B., Mermer, A., Bahar, H. (2005): An electrophoretic comparison of the venoms of a colubrid and various viperid snakes from Turkey and Cyprus, with some taxonomic and phylogenetic implications. *Zootaxa* 1038: 1-10.
- Atatür, M.K., Göçmen, B. (2001): Amphibians and Reptiles of Northern Cyprus [Kuzey Kıbrıs'ın Kurbağa ve Sürüngenleri] (English & Turkish). Ege Üniversitesi, Fen Fakültesi Kitaplar Serisi, No: 170, Ege Üniversitesi Basımevi, Bornova, İzmir.
- Bannikov, A.G., Darevsky, I.S., Ishchenko, V.G., Rustamov, A.K. and Szczerbak, N.N. (1977): *Opređelitel zemnovodnih i presmikayushchihysya fauni SSSR* [Guide to the amphibians and reptiles of the U.S.S.R. fauna], Prosveahchenie Press, Moscow.
- Baran, İ., Atatür, M.K. (1998): Turkish Herpetofauna (Amphibians and Reptiles). Republic of Turkey, Ministry of the Environment, Ankara.
- Baran, İ. (1976): Türkiye Yılanlarının Taksonomik Revizyonu ve Coğrafik Dağılımları [Taxonomic revision of Turkish Snakes and their geographic distributions]. TÜBTAK Yayınları, No: 309, T.B.A.G. Seri No: 9, Ankara.
- Başoğlu, M., Baran, İ. (1980): Türkiye Sürüngenleri. Kısım II: Yılanlar [The Reptiles of Turkey. Part II: Snakes]. Ege Üniversitesi, Fen Fakültesi, Kitaplar Serisi, No: 81, Ege Üniversitesi Matbaası, Bornova-İzmir.
- Budak, A., Göçmen, B. (2005): Herpetoloji. Ege Üniversitesi, Fen Fakültesi, Kitaplar Serisi No: 194, Bornova / İzmir.
- Billing, H., Schätti, B. (1984): Vorläufige Mitteilung zum Subspezies-Problem bei *Vipera lebetina* (Linnaeus 1758). *Salamandra* 20: 65-69.
- Böhme, W., Wiedl, H. (1994): Status and zoogeography of the herpetofauna of Cyprus with taxonomic and natural history notes on selected species genera (*Rana*, *Coluber*, *Natrix*, *Vipera*). *Amphibia and Reptilia. Zoology in the Middle East* 10: 31-52.

- Brodmann, P. (1987): Die Giftschlangen Europas und die Gattung *Vipera* in Afrika und Asien. Bern, Kümmerly-Frey, Bern.
- Chen, P.S. (1967): Separation of serum proteins in different amphibian species by polyacrylamide gel electrophoresis. *Experientia* 23: 483-485.
- Chernov, S.A. (1944): Taxonomic position of *Vipera lebetina* from Tajikistan. *Izv. Tadjh. Fil.AN SSSR. Zool. Parazitol.* 5:189-193.
- Chernov, S.A. (1959): Reptilia. The fauna of Tadj. SSR. Proc. Acad. Sci. Tadj. SSR, Stalinabad.
- Chikin, Y., Szczerbak, N.N. (1992): New subspecies of blunt-nosed viper, *Vipera lebetina chernovi* ssp. Nov. (Reptilia, Viperidae) from Middle Asia. *Vestn. Zool.* 6: 45-49.
- David, P., Ananjeva, N.B., Das, I., Golubev, M. (1999): Translation of the original description of *Vipera obtusa* with designation of a neotype. *Russian Journal of Herpetology* 6: 193-198.
- Dessauer, H.C., Fox, W. (1956): Characteristics electrophoretic pattern of orders of amphibia and reptilia. *Science* 124: 225-226.
- Eiselt, J., Baran, I. (1970): Ergebnisse zoologischer sammelreisen in der Türkei: Viperidae. *Annalen des Naturhistorischen Museums, Vienna*, 74: 357-369.
- Ferguson, A. (1980): Biochemical systematics and evolution. Blackie and Son 1-14.
- Göçmen, B., Tok, C.V., Kaya, U., Tosunoğlu, M. (1996): Kuzey Kıbrıs herpetofaunası hakkında bir ön çalışma raporu [A Preliminary study on the herpetofauna of northern Cyprus]. *Turkish Journal of Zoology* 20:161-176.
- Göçmen, B., Arıkan, H., Mermer, A., Langerwerf, B., Bahar, H. (2006): Morphological, Hemipenial, and Venom Electrophoresis Comparisons of the Levantine Viper, *Macrovipera lebetina* (Linnaeus, 1758), from Cyprus and Southern Anatolia. *Turkish Journal of Zoology* 30: 225-234.
- Joger, U. (1984): The venomous snakes of the Near and Middle East. pp. 1-112. In: Reichert, L. (ed.) Beihefte zum Tübinger Atlas des Vorderen Orients, Vol. 12, A. Naturwissenschaften, Weisbaden.
- Joger, U., Lenk, P., Baran, İ., Böhme, W., Ziegler, T., Heidrich, P., Wink, M. (1997): The phylogenetic position of *Vipera barani* and of *Vipera nikolskii* within the *Vipera berus* complex. pp.185-194. In: Böhme W., Bischoff, W., Ziegler, W.T. (eds.) *Herpetologia Bonnensis, Societas Europaea Herpetologica, Bonn.*
- Leviton, A.E., Anderson, S.C., Adler, K., Minton, S.A. (1992): Handbook to Middle East Amphibians and Reptiles. Society for the Study of Amphibians and Reptiles, Oxford (Ohio), U.S.A.
- MacLean, G.S., Lee, S.K., Wilson, K.F. (1973): A simple method of obtaining blood from lizards. *Copeia* 2: 338-339.
- Martin, W.C.L. (1838): On some snakes collected during the Euphrates expedition. *Proc. Zool. Soc.* 1838: 81-84.
- Mertens, R. (1952): Amphibien und reptilien aus der Türkei. *İstanbul Üniversitesi Fen Fakültesi Mecmuası Seri B, İstanbul*, 17:41-75.
- Mulder, J. (1995): Herpetological observations in Turkey (1987-1995). *Deinsa* 2: 51-66.
- Nilson, G., Andrén, C., Flärdh, B. (1988): Die vipern der Türkei. *Salamandra* 24: 215-247.
- Nilson, G., Andrén, C. (1988): *Vipera lebetina transmediterranea*, a new subspecies of viper from North Africa, with remarks on the taxonomy of *Vipera lebetina* and *Vipera mauritanica*. *Bonn. Zool. Beitr.* 39: 371-379
- Osenegg, K. (1989): Die amphibien und reptilien der Insel Zypern. University of Bonn (The thesis of M. Sc.), Bonn.
- Schätti, B., Sigg, H. (1989): Die herpetofauna der Insel Zypern. Teil 2. Schildkröten, Echsen und Schlangen. *Herpetofauna* 11(62): 15-26.
- Schleich, H. H., Kästle, W., Kabish, K. (1996): Amphibians and Reptiles of North Africa. Koeltz Scientific Books, Germany.
- Terentjev, P., Chernov, S. A. (1949): Key to amphibians and reptiles [in Russian]. Israel Program for Scientific Translation, Jarusalem, 1965.
- Tok, C.V., Cihan, D., Ayaz, D. (2002): A new record of *Macrovipera lebetina obtusa* (Viperidae) from South-Eastern Anatolia. *Zoology in the Middle East* 25:23-26.

Submitted: 27 August 2007
/ Accepted: 29 September 2007

Corresponding Editor: I. Sas
English Language Editor: A.P. Pernetta